

Three-dimensional magnetized and rotating hot plasma equilibrium and stability in a gravitational field

Peter J. Catto¹, Sergei I. Krashennnikov², Istvan Pusztai³

¹ *Plasma Science and Fusion Center, MIT, Cambridge, MA 02139, USA*

² *University of California at San Diego, La Jolla, CA 92093, USA*

³ *Department of Applied Physics, Chalmers U. of Technology, 41296 Gothenburg, Sweden*

Abstract

We present analytic solutions for three-dimensional magnetized axisymmetric equilibria confining rotating hot plasma in a gravitational field. Our solution to the full Grad-Shafranov equation can exhibit strong equatorial plane localization of the plasma density and current, resulting in disk equilibria for the plasma density. Unlike in [1], we find a toroidal magnetic field is necessary to find an equilibrium in the presence of gravity for most cases of interest. We expect our results to provide impetus to re-investigate magneto-rotational stability [2,3].

Introduction: Fully self-consistent three dimensional global equilibrium of hot, rotating plasma confined by gravity and magnetic field are of interest for astrophysical and space plasma applications, but have proven difficult to find. Previous magnetohydrodynamic (MHD) models assume strict incompressibility with constant density [4], assume the density is a flux function [5], ignore the frozen in constraint [6], and/or require poloidal flow [5-7] or an adiabatic equation of state [8,9], which are not allowed kinetically. To satisfy constraints imposed on a drifting Maxwellian by the Fokker-Planck equation, only toroidal flow is allowed and the temperature must be a flux function [1,10-12]. Moreover, in the presence of rotation and gravity the density must be allowed to vary poloidally as well as radially since strict Keplerian motion is only possible at the equatorial plane [1]. It is anticipated that the equilibria we find will be useful in setting up global simulations to investigate magneto-rotational stability [2,3] in accretion disks and help shed light on the detailed mechanisms by which mass accretion occurs at a black hole as momentum is transported outward.

Grad-Shafranov equation: The flux surfaces for an axisymmetric equilibrium must satisfy a Grad-Shafranov equation that we find from Ampere's law and pressure balance by taking the magnetic field to be given by

$$\vec{B} = I\nabla\zeta + \nabla\psi \times \nabla\zeta, \quad (1)$$

where ζ is the toroidal angle, ψ is the poloidal flux function, $I = RB_T$ with B_T the toroidal magnetic field and R the cylindrical radius from the axis of symmetry. To satisfy the kinetic constraints [10-12] the velocity $\vec{V}$ must be toroidal

$$\vec{V} = \Omega R^2 \nabla \zeta, \quad (2)$$

with $\Omega = c d\Phi/d\psi$ the toroidal rotation frequency. The magnetic field is frozen into the flow so that $c\nabla\Phi = \vec{V} \times \vec{B}$ gives $\vec{B} \cdot \nabla\Phi = 0$, requiring the electrostatic potential Φ to be a flux function to lowest order. In addition, we write the gravitational potential G as

$$G = -G_0 M_0 / r \quad (3)$$

with G_0 the gravitational constant, r the spherical radius, and M_0 the mass of the astrophysical body that is assumed to be a compact source centered at $r = 0$, such that $R = r \sin\vartheta$ with ϑ the angle from the axis of symmetry. Our spherical and cylindrical coordinates are defined to satisfy $r\nabla\vartheta = R\nabla\zeta \times \nabla r$ and $\nabla z = R\nabla R \times \nabla\zeta$.

Conservation of momentum requires

$$c^{-1} \vec{J} \times \vec{B} = Mn(\nabla G - \Omega^2 R \nabla R) + \nabla p. \quad (4)$$

The parallel component requires that the ion density n depend on poloidal angle

$$n = n(\psi, \vartheta) = \eta(\psi) e^{\kappa(\psi, \vartheta)} = \eta(\psi) e^{\kappa(\psi, \vartheta)}, \quad (5)$$

where the density and normalizing or "pseudo" density η are related via the generalized Maxwell-Boltzmann exponential factor κ that retains the poloidal dependence due to the centrifugal and gravitational potential,

$$4T\kappa/M = \Omega^2 R^2 - 2G, \quad (6)$$

with $T = T(\psi)$ and $p = n(T_i + ZT_e) = 2nT$ in a quasi-neutral plasma ($Zn = n_e$ = electron density) of ion charge number Z and ion and electron temperatures T_i and T_e . The toroidal momentum balance requires the current density, $\vec{J}$, across a flux surface vanish, $\vec{J} \cdot \nabla\psi = 0$, and then Ampere's law gives $I = I(\psi)$. Ampere's law also gives $4\pi c^{-1} \vec{J} \cdot \nabla\vartheta = \vec{B} \cdot \nabla dI/d\psi$. Then the current density can be conveniently decomposed into toroidal and parallel components by writing it as $\vec{J} = R\vec{J}_* \nabla\zeta + (c/4\pi) dI/d\psi \vec{B}$, allowing us to obtain the useful relation $\vec{J} \cdot \vec{B} = I\vec{J} \cdot \nabla\zeta + (cB_p^2/4\pi R^2) dI/d\psi$, with $\vec{B}_p = \nabla\psi \times \nabla\zeta$ the poloidal magnetic field and $B_p^2 = R^{-2} |\nabla\psi|^2$. Combining this result with the $\nabla\psi$ component of force balance, $R^2 B^2 \vec{J} \cdot \nabla\zeta = I\vec{J} \cdot \vec{B} + c\nabla\psi \cdot [\nabla p + Mn(\nabla G - \Omega^2 R \nabla R)]$, and the toroidal component of Ampere's law yields the Grad-Shafranov equation [13]

$$\nabla \cdot (R^{-2} \nabla\psi) = -IR^{-2} dI/d\psi - 4\pi R^{-2} B_p^2 \nabla\psi \cdot [\nabla p + Mn(\nabla G - \Omega^2 R \nabla R)]. \quad (7)$$

Separable form: To find a separable form for the Grad-Shafranov equation we assume

$$\psi = \psi_0 H(\mu) (R_0/r)^\alpha, \quad (8)$$

where $\mu = \cos\vartheta$ and our normalization is $H(\mu=0) = 1$. The vacuum limit $H = 1 - \mu^2$ recovers a homogeneous magnetic field for $\alpha = -2$ and a point dipole solution if $\alpha = 1$, and ψ_0 is a constant reference value at the reference location R_0 (reference values are defined at the equatorial plane $\mu = 0$ and denoted by a subscript "o"). We obtain an ordinary differential equation for H by assuming

$$\Omega^2 = \Omega_0^2 (\psi / \psi_0)^{3/\alpha} \propto 1 / r^3, \quad (9)$$

$$T = T_0 (\psi / \psi_0)^{1/\alpha}, \quad (10)$$

$$I = I_0 (\psi / \psi_0)^{1+1/\alpha}, \quad (11)$$

$$n = n_0 (\psi / \psi_0)^{2+3/\alpha} e^{\kappa(\mu)}, \quad (12)$$

with $n_0 = n(\psi = \psi_0, \mu = 0)$. Defining the positive constants

$$g = \frac{8\pi G_0 M_0 M n_0}{B_{p0}^2 R_0} = \frac{8\pi G_0 M_0 M n}{B_p^2 r} \Big|_{\mu=0}, \quad (13)$$

$$\omega^2 = \frac{4\pi M n_0 \Omega_0^2 R_0^2}{B_{p0}^2} = \frac{4\pi M n \Omega^2 r^2}{B_p^2} \Big|_{\mu=0}, \quad (14)$$

$$\beta = \frac{16\pi n_0 T_0}{B_{p0}^2} = \frac{8\pi p}{B_p^2} \Big|_{\mu=0}, \quad (15)$$

and

$$b^2 = \frac{I_0^2}{R_0^2 B_{p0}^2} = \frac{I^2}{R^2 B_p^2} \Big|_{\mu=0} = \frac{B_T^2}{B_p^2} \Big|_{\mu=0}, \quad (16)$$

an ordinary second order nonlinear differential equation for H is obtained from (7) [13]:

$$\frac{d^2 H}{d\mu^2} + \frac{\alpha(\alpha+1)}{1-\mu^2} H = \alpha \left[\frac{g}{2} H^{-1/\alpha} - \omega^2 (1-\mu^2) H^{2/\alpha} - (\alpha+2)\beta - \frac{(\alpha+1)b^2}{(1-\mu^2)H^{2/\alpha}} \right] H^{1+4/\alpha} e^{\kappa}, \quad (17)$$

where

$$\kappa = -(g/\beta)(1-H^{-1/\alpha}) - (\omega^2/\beta)[1-(1-\mu^2)H^{2/\alpha}], \quad (18)$$

Solution technique: Solutions to this Grad-Shafranov equation can be found by the techniques illustrated in [1] and references therein, however, some care is needed. The solution H must be up-down symmetric and monotonically decreasing from unity until the poloidal magnetic field vanishes at the axis of symmetry ($1 \geq H \geq 0$). This behavior seems to require $d^2 H / d\mu^2 < 0$ for all μ . Consequently, we desire $g - (\alpha+1)(2b^2+1) > 2\omega^2 + 2(\alpha+2)\beta$ at $\mu = 0$. Moreover, gravity forces us to assume $\alpha < 0$ to avoid singular behavior as $H \rightarrow 0$, with rotation requiring $H \leq (1-\mu^2)^{-\alpha/2}$ to avoid exponential growth at the poles. An exact solution,

$H=(1-\mu^2)^{-\alpha/2}$, exists if $G = 0$ [1] when $\alpha = -[2(\beta+1)+\omega^2+b^2]/(\beta+1+b^2)$. The poloidal magnetic field decreases with radius if $\alpha > -2$ and the toroidal magnetic field remains finite at the poles if $\alpha < -1$ so we expect solutions with $-1 > \alpha \geq -2$ are of most interest in the presence of gravity. The $g = 0$ solution suggests maintaining $d^2H/d\mu^2 < 0$ as $\mu^2 \rightarrow 1$ requires $[(\alpha+1)b^2 > \omega^2 + (\alpha+2)\beta]$, a condition that is needed even when g is retained since $H^{-1/\alpha} \rightarrow 0$. This restriction implies a toroidal magnetic field is needed to obtain a fully self-consistent gravitational equilibrium. However, the procedures of [1] remain valid for obtaining solutions for $-1 > \alpha \geq -2$ with b^2 retained. Consequently, it is possible to find plasma disk equilibrium solutions to the Grad-Shafranov equation (17) and (18) that predict the rotation frequency bound, and the conditions for a plasma disk as well as its thickness.

Magneto-rotational instability (MRI) without gravity: In the absence of gravity the toroidal magnetic field can be neglected. In this case $\alpha < -2$, but the MRI calculation is more tractable [2]. The equilibrium associated with our exact solution for H is unstable if $\beta \gg 1$ and $\Omega^2 R^2 < 10T/M$ in the incompressible limit (roughly in agreement with [2]), and when $\beta < 3/2$ and $\Omega^2 R^2 \gg 2T/M$ in the compressible limit not considered in [2]. In the presence of gravity [3] the toroidal magnetic field must be retained to investigate MRI instability.

Work supported by the US Department of Energy grants DE-FG02-91ER-54109 at MIT and DE-FG02-04ER54739 at UCSD & by an International Career Grant from Vetenskapsrådet.

- [1] CATTO, P. J. & KRASHENINNIKOV S. I. 2015 A rotating and magnetized three-dimensional hot plasma equilibrium in a gravitating field. *J. Plasma Physics* **81**, 105810301.
- [2] VELIKHOV, E. P. 1959 Stability of an ideally conducting liquid flowing between cylinders rotating in a magnetic field. *Sov. Phys. JETP* **36**, 995-998 (1959).
- [3] BALBUS, S. A. & HAWLEY, J. F. 1991 A powerful local shear instability in weakly magnetized disks. I. Linear analysis, *Astrophys. J.* **376**, 214-222.
- [4] CHANDRASEKHAR, S. 1956 Axisymmetric magnetic fields and fluid motions. *ApJ* **124**, 232-243.
- [5] THROUMOULOPOULOS, G. N. & TASSO, H. 2001 Axisymmetric equilibria of a gravitating plasma with incompressible flows. *Geophys. Astroph. Fluid Dynamics* **94**, 249-262.
- [6] PRASANNA, A. R., TRIPATHY, S. C. & DAS, A. C. 1989 Equilibrium structure for a plasma magnetosphere around compact objects. *J. Astrophys. Astr.* **10**, 21-34.
- [7] MCCLEMENTS, K. C. & THYAGARAYA, A. 2001 Azimuthally symmetric magnetohydrodynamic and two-fluid equilibria with arbitrary flows. *Mon. Not. Roy. Astr. Soc.* **323**, 733-742.
- [8] LOVELACE, R. V. E., MEHANIAN, C., MOBARRY, C. M. & SULKANEN, M. E. 1986 Theory of axisymmetric magnetohydrodynamic flows: disks. *ApJS* **62**, 1-37.
- [9] OGILVIE, G. I. 1997 The equilibrium of a differentially rotating disc containing a poloidal magnetic field. *Mon. Not. R. Astron. Soc.* **288**, 63-77.
- [10] HINTON, F. L. & WONG, S. K. 1985 Neoclassical ion transport in rotating axisymmetric plasmas. *Phys. Fluids* **28**, 3082-3098.
- [11] CATTO, P. J., BERNSTEIN, I. B. & TESSAROTTO, M. 1987 Ion transport in toroidally rotating tokamak plasmas. *Phys. Fluids* **30**, 2784-2795.
- [12] HELANDER, P. 2014 Theory of plasma confinement in non-axisymmetric magnetic fields. *Rep. Prog. Phys.* **77**, 087001-35.
- [13] KRASHENINNIKOV, S. I., CATTO, P. J. & HAZELTINE, R. D. 2000 Plasma equilibria in dipolar magnetic configurations. *Phys. Plasmas* **7**, 1831-1838.